



DESCRIPTION

Glycosade is a hydrothermally processed high amylopectin starch.

INDICATIONS

For use in the dietary management of Glycogen Storage Disease (GSD) where the use of a long acting starch is indicated.

Suitable from 5 years of age.

NUTRITION INFORMATION

| | | per 100g | per 60g packet |
|-----------------|------|----------|----------------|
| Calories | kcal | 356 | 214 |
| Protein | g | 0.5 | 0.3 |
| Carbohydrate | g | 88 | 53 |
| Sugars | g | 0 | 0 |
| Fat | g | 0.14 | 0.08 |
| Saturates | g | 0.09 | 0.05 |
| Fiber | g | 0 | 0 |
| Minerals | | | |
| Sodium | mg | 20 | 12 |
| | mmol | 0.9 | 0.5 |

Medical food intended for use under medical supervision.

DOSAGE AND ADMINISTRATION

To be determined by the clinician or dietitian following appropriate assessment (a fasting Glycosade load is recommended) and is dependent on the age, body weight and medical condition of the patient.

Each 60g packet of Glycosade has an equivalent carbohydrate content to approximately 55g of uncooked cornstarch.

PREPARATION GUIDELINES

To be determined by the clinician or dietitian.

Glycosade taken as a drink:

1. Measure Glycosade dose into a cup with a lid.
2. Add the liquid of your choice. Typically a packet of Glycosade can be added to 100ml of cold water or other permitted fluids.
3. Close the lid tightly and shake well for 10 seconds or until powder is dissolved.
4. Drink Glycosade fresh every time.

IMPORTANT NOTICE

Must only be used in patients with proven GSD where the use of a long acting starch is indicated.

Suitable from 5 years of age in the USA.
Suitable from 2 years of age in Canada.

Not for use as a sole source of nutrition.

Not for intravenous use.

STORAGE

Unopened: Store in a cool, dry place.

Once opened: Ideally packets should be used immediately after opening.

Any unused powder in an open packet should be kept in an airtight container or sealed with a clip and used within 24 hours.

PACK SIZE

30 x 60g (2.1 oz) packets.

WEIGHT

1.8kg (3.97 lb)

SHELF LIFE

12 months.



Innovation in Nutrition

Manufactured in the UK for:
VitaFlo USA LLC, 211 North Union Street,
Suite 100, Alexandria, VA 22314.
Tel: 1-888-VITAFLO
Fax: 631-693-2002
www.VitaFloUSA.com
E-mail: vitaflo@vitaflousa.com

VitaFlo and Glycosade are Reg. Trademarks of Société des produits Nestlé S.A., Vevey, Switzerland.

Code: January 2014